Climate Change and Human Health Literature Portal



The impact of excess heat events in Maricopa County, Arizona: 2000--2005

Author(s): Yip FY, Flanders WD, Wolkin A, Engelthaler D, Humble W, Neri A, Lewis L,

Backer L, Rubin C

Year: 2008

Journal: International Journal of Biometeorology. 52 (8): 765-772

Abstract:

Exposure to excess heat is preventable yet it is the primary weather-related cause of mortality in the United States. In the Southwest United States, high temperatures are common and indoor environments often have cooling devices. In summer 2005, Maricopa County, Arizona experienced a 182% increase in reported heat-related deaths in comparison to 2000--2004. We examined at-risk populations and excess mortality. We characterized heat-related deaths using descriptive and multivariate time-series analyses of county vital record data from June-September 2000--2005. Dose-response relationships for heat-related mortality and heat index were evaluated using linear and guadratic splines. From June-September, 2000--2005, 136 heat-related deaths (0.68 per 100,000) were reported; 49 (36%) occurred in 2005. In July 2005, a 14-day heat wave resulted in 28 (57%) reported deaths--a 102% increase in comparison to the same time period in 2000--2004. Decedent demographics in 2005 did not differ from previous years. The mean age of all 136 deaths was 56 years (range: 7-92 years). Of those with discernable reported injury locations, 62 (66%) were identified outdoors. Forty-eight (77%) decedents identified outdoors were < 5 years; conversely, 26 (82%) decedents who were found indoors were > or Euro Surveillance (Bulletin Europeen Sur Les Maladies Transmissibles; European Communicable Disease Bulletin) 65 years. A 6% (95% CI: 1.00-1.13) increase in mortality risk was observed for each degree (F) increase in heat index. Excess heat impacted a younger population in Maricopa County and many deaths occurred outdoors. Consecutive days of heat exposure--even among a heat-acclimated population--can increase mortality risk. Public health response activities guided by locally obtained data will better target those at risk.

Source: http://dx.doi.org/10.1007/s00484-008-0169-0

Resource Description

Communication: M

resource focus on research or methods on how to communicate or frame issues on climate change; surveys of attitudes, knowledge, beliefs about climate change

A focus of content

Other Communication Audience: Local governments

Early Warning System:

resource focus on systems used to warn populations of high temperatures, extreme weather, or other elements of climate change to prevent harm to health

Climate Change and Human Health Literature Portal

A focus of content
Exposure:
weather or climate related pathway by which climate change affects health
Temperature
Temperature: Extreme Heat
Geographic Feature: 🛚
resource focuses on specific type of geography
Desert
Geographic Location: 🖫
resource focuses on specific location
United States
Health Impact: ☑
specification of health effect or disease related to climate change exposure
Injury
Mitigation/Adaptation: ₪
mitigation or adaptation strategy is a focus of resource
Adaptation
Population of Concern: A focus of content
Population of Concern: ₪
populations at particular risk or vulnerability to climate change impacts
Elderly
Resource Type:
format or standard characteristic of resource
Research Article
Timescale: ™
time period studied
Time Scale Unspecified
Vulnerability/Impact Assessment: ☑
resource focus on process of identifying, quantifying, and prioritizing vulnerabilities in a system
A focus of content